

PRECISION

Intra-assay Precision (Precision within an assay): CV%<8%


Three samples of known concentration were tested twenty times on one plate to assess.

Inter-assay Precision (Precision between assays): CV%<10%

Three samples of known concentration were tested in twenty assays to assess.

TYPICAL DATA

These standard curves are provided for demonstration only. A standard curve should be generated for each set of samples assayed.



ng/ml	OD1	OD2	Average	Corrected
6000	2.755	2.654	2.705	2.615
3000	2.028	1.917	1.973	1.883
1500	1.195	1.174	1.185	1.095
750	0.569	0.558	0.564	0.474
375	0.322	0.312	0.317	0.227
187.5	0.254	0.243	0.249	0.159
93.75	0.148	0.144	0.146	0.056
0	0.090	0.089	0.090	

LOD

23.4 ng/ml

LINEARITY

To assess the linearity of the assay, samples were spiked with high concentrations of guinea pig IgE in various matrices and diluted with the Sample Diluent to produce samples with values within the dynamic range of the assay.

	Sample	Serum(n=4)
1:1	Average %	89
	Range %	82-96
1:2	Average %	94
	Range %	91-97
1:4	Average %	103
	Range %	96-107
1:8	Average %	98
	Range %	92-102

RECOVERY

The recovery of guinea pig IgE spiked to levels throughout the range of the assay in various matrices was evaluated. Samples were diluted prior to assay as directed in the Sample Preparation section.

Sample Type	Average % Recovery	Range
Serum (n=5)	89	82-96
EDTA plasma (n=4)	106	102-110